

DOLODONOV, S., podpolkovnik

Radiation and chemical reconnaissance. Voen. vest. 41  
no.5:50-53 My '61. (MIRA 14:8)  
(Stream crossing, Military) (Chemical warfare)  
(Radioactive fallout)

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000410830003-2

DOLODONOV, S., podpolkovnik

Under reliable protection. Voen. vest. 42 no.8:48-51 Ag  
'62.

(Chemical warfare--Safety measures)

(MIRA 15:7)

APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000410830003-2"

S/018/63/000/002/001/001  
A004/A126

AUTHORS: Kolyada, I., Dolodonov, S., Lieutenant-Colonels

TITLE: Efficient protection from radioactive substances and toxic agents

PERIODICAL: Voyenny vestnik, no. 2, 1963, 54 - 55

TEXT: The authors describe ways and means for an efficient protection from radioactive substances, toxic agents and war gas in winter. They point out that after snow-fall and by snow-storms radioactive fallout may be carried by snow and storm to areas where a contamination was not to be expected, and that radiation monitors do not operate accurately at low temperatures. They enumerate the various measures and monitoring methods that are to be applied by the radiation detection teams, and describe the behavior and protective clothing and means of an attacking battalion in a contaminated zone, as well as the various disinfection measures to be taken to make sure that all radioactive substance and toxic agents are removed.

Card 1/1

DOLOGOFOLOV, V.I., inzh; DOLOGOPOLAV, L.N., inzh; PETROVA, N.G., inzh;  
MILYAYEVA, T.I., inzh.

Electroluminescent panels. Svetotekhnika 8 no.11:24-26 N 1962.

(MIRA 15:10)

1. Vsesoyuznyy svetotekhnicheskiy institut.  
(Luminiscence) (Electric apparatus and appliances)

DOLOGOPOLOV, V.I., inzh; DOLOGOPOLAV, L.N., inzh; PETROVA, N.G., inzh;  
MILYAYEVA, T.I., inzh.

Electroluminescent panels. Svetotekhnika 8 no.11:24-26 N. '62.

(MIRA 15:10)

1. Vsesoyuznyy svetotekhnicheskiy institut.  
(Luminescence) (Electric apparatus and appliances)

V. N.; DAYON, M. I.; DEVISHEV, M. I.; DOLOGOSHEYN, B.A.; KLIMANOVA, Z.  
S. I.; SHMEL'VA, A. P.

Discharge Track-Detector Chamber Investigation of Characteristics of some  
Dark Chambers.

submitted for the Intl. Conf. on Cosmic Rays (IUPAP), Jaipur India,  
Dec 1963.

1. DOLOGOSHOV V.
2. USSR (600)
4. Shrubs
7. Ripening and fall of fruit and seeds from trees and bushes. Les.khoz. 5  
no. 12, 1952.
9. Monthly List of Russian Acquisitions, Library of Congress, April 1953, unclass

DOLOKO, A.L., otvetstvennyy red.

[Kazan railroad on the fortieth anniversary of the Great October  
Socialist Revolution; an account of its history and economy]  
Kazanskaya zheleznaia doroga k 40-ii godovshchine Velikoi Oktiabr'-  
skoi sotsialisticheskoi revoliutsii; istoriko-ekonomicheskii  
ocherk. Kazan', 1957. 39 p.  
(MIRA 11:6)

1. Russia (1923- U.S.S.R.) Kazanskaya zheleznaya doroga.  
Dorozhnoe nauchno-tehnicheskoye obshchestvo. 2. Glavnyy  
inzhener Kazanskoy zheleznoy dorogi.  
(Railroads)

DOLOMAKIN, P., mayster po kladtsi kil'tsevikh pechey

Laying annular kilns. Sil'. bud. 9 no.5:21-22 My '59.  
(MIRA 13:3)  
(Kilns)

L 515-66 EWT(1)/T IJP(c) 3  
ACC NR. AP5025340 SOURCE CODE: UR/0126/65/020/003/0474/0475

AUTHOR: Mezentseva, N. L.; Mikhayl'eva, N. P.; Starostina, L. S.; Dolomanov, L. A.

ORG: Institute of Crystallography, AN SSSR (Institut Kristallografii AN SSSR)

TITLE: Growing and refining of yttrium single crystals

SOURCE: Fizika metallov i metallovedeniye, v. 20, no. 3, 1965, 474-475

TOPIC TAGS: yttrium, yttrium zone refining, zone refining, yttrium single crystal, single crystal growth

ABSTRACT: Experiments have been made to combine vacuum crucibleless zone refining of yttrium with simultaneous growing of yttrium single crystals. Yttrium ingots 80 mm long were subjected to crucibleless zone refining during which yttrium single crystals up to 1 mm long and 5 mm in diameter were successfully grown. A polymorphic transformation in yttrium at 1475°C made it impossible to obtain large single crystals. Zone refining decreased the content of iron, copper, and calcium in the ingots by 27, 20, and 10 times, respectively. The oxygen content decreased by only 2.4 times, presumably because of the formation of stable yttrium oxides, whose distribution coefficient is close to 1. For more effective refining of the growing single crystals, electric current was passed through the ingot simultaneously with the zone fusion. After 8-hr processing at a current density of 5 amp/mm<sup>2</sup>, the oxygen content in the ingot anode portion, simultaneously zone fused, was three times lower than that near the cathode portion. Orig. art. has 1 figure. [MS]

Card 1/2 UDC: 669.794.172

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000410830003-2

L 5115-66

ACC NR: AP5025340

SUB CODE: S8/ SUBM DATE: 2000ct64/ ORIG REF: 002/ OTH REF: 001/ ATD PRESS: 4/33

PC  
Card 2/2

APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000410830003-2"

26.2/14

39787  
S/147/62/000/002/015/020  
E191/E535

AUTHOR: Dolomanov, Ye.G.

TITLE: Inversion phenomena in real gases in supersonic flow

PERIODICAL: Izvestiya vysshikh uchebnykh zavedeniy, Aviatsionnaya tekhnika, no.2, 1962, 124-129

TEXT: In the solution of many technical problems, the magnitude of the stagnation temperature after a compression shock must be known. For an ideal gas, the stagnation temperature after the shock is assumed equal to that before the shock. In modern aircraft, the stagnation quantities (pressure and temperature) before the compression shock are so high that relationships applicable to real gases should be used. The change of enthalpy in the stagnation process is formulated; the total enthalpy remains constant when a real gas traverses a compression shock front. From these statements, the change in the stagnation temperature is derived in terms of the initial flow variables. A discussion of the expression so obtained making use of temperature-volume diagrams of typical real gases shows that a compression shock can either increase, or decrease, or have no effect on, the stagnation temperature. When the stagnation temperature remains

Card 1/2

Inversion phenomena in real gases ... S/147/62/000/002/015/020  
E191/E535

constant the conditions are referred to as inversion point. There is an inversion line in the temperature/volume diagram joining all inversion points. The change in the stagnation temperature of a real gas through a compression shock is proportional to its change of total pressure. The proportionality factor, the "differential shock effect", is dependent on the initial flow variables and the specific heat. This effect has the same physical significance as the Joule-Thompson effect. The relation of the shock effect to other properties of the gas is discussed. When a gas is more compressible than an ideal gas the stagnation temperature drops through the shock; if less compressible, it increases with the shock. There are 3 figures.

SUBMITTED: October 13, 1961

Card 2/2

L 13590-63 EPR/EPA(b)/EMT(1)/EDS AFFTC/ASD Pe-4/Pd-4 WH  
ACCESSION NR: AP3004723 8/0147/63/000/002/0065/0069

AUTHOR: Dolmanov, Ye. G.

61

TITLE: On the calculation of air parameters behind a normal shock with stagnation-temperature inversion taken into account

SOURCE: IVUZ. Aviats. tekhnika, no. 2, 1963, 65-69

TOPIC (ACOS): stagnation temperature, stagnation-temperature inversion, normal shock, stagnation region, flow, shock front, shock effect

ABSTRACT: An analysis of the conclusions made by the author in his previous work (IVUZ., Aviatsionnaya tekhnika, no. 2, 1962) concerning the existence, at the shock, of stagnation-temperature inversion is presented. The flow parameters in the region of the stagnation point are calculated, taking the effect of temperature inversion into account. A graph-analytical method is used for calculation of the gasdynamic and thermodynamic parameters of air (see Fig. 1 of Enclosure); the effects of ionization, dissociation, NO molecule formation, and variable specific heat are considered. The importance of temperature inversion, called shock effect, is demonstrated, and a comparison of the observed and theoretical results is presented in a table (see Enclosure). It is concluded that shock effect is

Cord 1/M2

L 13590-63  
ACCESSION NR: AP3004723

present in all gases, and that dissociation, ionisation, NO molecule formation, variable specific heat, and shock effect act in the same direction, that is, they decrease stagnation temperature. It is also concluded that the graph-analytical method is much simpler for calculating gasdynamic and thermodynamic parameters than the mathematical method. Orig. art. has: 3 figures, 15 formulas, and 1 table.

ASSOCIATION: none

SUBMITTED: 23Nov62

DATE ACQ: 06Sep63

ENCL: 02

SUB CNDIS: AI

NO REP Sov: 005

OTHER: 001

Card 2/42

DOLOMANDVA, N.P.

Evaluation of the ampic method for determination of sodium chloride in cheese. Z. B. Kristall, N. P. Dolmanova, I. I. Golubova, and R. I. Merenchenko. *Sbornik Nauchno-tekhnicheskikh Rabot Nauchno-Issledovatel'skogo Instituta po Molekulyarnoi Promst. 1934, No. 3, 21-31; Referat. Akad. Nauk SSSR po Khim. i Khim. Tekhnologii, 1934, No. 43400.*—The argentometric and mercurimetric methods are compared. In the argentometric method the losses of Cl in ashing are  $\leq 23\%$ . The mercurimetric method is recommended as the ampic method. Treat 1 g. of cheese with 10 ml. of 1*N* KOH while heating, cool, acidify with HNO<sub>3</sub> to a concn. of 1*N*, and titrate with 1*N* HgNO<sub>3</sub>, adding toward the end diphenylcarbazole. At control boil the cheese with KMnO<sub>4</sub> in a weakly alk. soln. until a stable violet color is obtained. Acidify with excess HNO<sub>3</sub>, treat with H<sub>2</sub>O<sub>2</sub> to disappearance of pink color, and titrate with HgNO<sub>3</sub>. The proposed analysis takes 40 min. and that with wet oxidation 1.5 hrs. The results compare well.

M. J. Koch

(3)

DOLJANOV, re. 1., and GRIGOR'YEV, I. F.

new data on crystallochemistry and typomorphic features of cassiterite of various origins. Study min. mus. no 3, 1951.

DOLOMANOVA, E.I.

USSR

Crystalliochemical and typomorphitic properties of cassiterite of different genesis. IV. N. Grigor'ev and R. I. Dolomanova. *Izv. Akad. Nauk S.S.R. Min. No. 3, 73-92 (1951).*—The most important accessory elements occurring in cassiterite are Nb, Ta, W, Fe, Mn, Zr, and Ti. These elements do not replace Sn by isomorphism but form independent mineral assem., namely of columbite, wolframite, zircon, and rutile, most frequently in regular intergrowths. They form crystallites of only 1  $\mu$  size and smaller, i.e. in such a degree of dispersity that they act as allochromic pigments of cassiterite. Columbite, wolframite, and hematite are only found in the dark-colored zones, and "hour-glass" structures of cassiterite crystals but not in the colorless zone. Dark-colored cassiterite is

(Due 2-1)

GP

①

therefore paramagnetic, while pure  $\text{SnO}_2$  is diamagnetic. The colorless zones are practically pure, and include only distinct crystallites of quartz or silicates. Among the accessory elements, Nb and W indicate to a certain degree the paragenetic and geochem. conditions under which the Sn deposits have been formed, or the geochem. character of the geological province in which they occur. Ag and Sb are also of this type; they are exclusively restricted to cassiterites which have been formed in sulfide mineral deposits. Typically bipyramidal crystals of cassiterite are characteristic for the metasomatic granite formation, which is associated with pegmatites and granites or cryst. schists, which have undergone a rapid temp. decrease on narrow space conditions of crystal growth. This bipyramidal habit of cassiterite is always specifically correlated with Nb (besides Fe and Mn) as "indicator" element. On the other hand, the prismatic crystal habit of cassiterite occurring in quartz-ore veins is always correlated with W as indicator element (besides Fe and Mn in wolframite and molybdenite). Details are given on biaxial cassiterite which is always rich in contaminations, while colorless  $\text{SnO}_2$  is unusual. Columbite, wolframite, etc., do not form solid solutions with cassiterite, but crystallize independently. This agrees with the results of Neuhäus and Noll (C.A. 44, 4377d). W. Bittel

DOLOMANOVA, Ye.I.

Gekkaitse, a new water-containing calcium aluminum fluorite mineral, by P. Grigor'ev and Ye.I. Dolomanova, Drude Mineralog. Akad. Nauk SSSR, No. 1, "93-01851."—A white mineral of kaolinite-like habit was observed in the oxidation zone of deposits in Transbaikalia region with the formula  $\text{Ca}_2\text{Al}_2(\text{OH})_2\text{Si}_2\text{O}_5$ , called gekkaitse. The rocks in which the mineral has formed are hornfels-like metamorphic sediments (erratically clay-slates and sandstones) which were changed by the hydrothermal veins, ascending fissure cracks and veins swarming through the country rocks. Topaz, staurolite, amosite, fluorapatite and sulfides are primary minerals; secondary minerals are halloysite, montmorillonite, and seattleite. The chem. analysis of the new mineral (with  $\text{Al}_2\text{O}_3$  44.42%;  $\text{CaO}$  14.30%;  $\text{H}_2\text{O}$  + 15.39%;  $\text{F}$  37.10%) and its x-ray powder diagram shows its independent character, in the group of gekkaitse,  $\text{Ca}[\text{Al}(\text{OH})_4]_2$ , para-gekkaitse,  $\text{Ca}[\text{Al}(\text{F},\text{OH})_4]_2\text{H}_2\text{O}$ , and belyanitite,  $\text{Ca}[\text{Al}(\text{P},\text{OH})_4]_2\text{H}_2\text{O}$ . Gekkaitse forms dense aggregates with a birefringence of less than 0.000, about 1.189. The spectral analysis detected only weak lines of Fe, Cr, As, Mn, Ni, and traces of Pb, Zn, Na, Co, Ti. The differential thermal analysis is characterized by strong endothermic peaks at 380°, 490°, 1320°, and a slight exothermic peak at 730°. Gekkaitse is explained as a supergene replacement product of topaz and fluorite.

*DOLOMANOVA, K.*

/ Stannite, and the products of its decomposition. lv. F. G.P.

Grigor'ev and L. I. Dolomanova. *Trudy Mineralog.*

*Muzh. Akad. Nauk SSSR No. 4, 74-76 (1982).* --Pre-

vailingly fine-cryst. aggregates of stannite are observed, assoc. with chalcopyrite, arsenopyrite, and sphalerite in Karella, the Caucasus, Transbaikalia, and the coast districts of northeastern Siberia. Single crystals with the pseudoregular forms  $\{11\}$  and  $\{001\}$  are rare, occurring in druses in quartz veins with chalcopyrite and arsenopyrite. A Zn-contg. stannite has a somewhat different x-ray structure than common stannite. The mineral is often changed on its surface to an olive-green "hydrostannite" (about  $\text{Fe}_2\text{O}_3 \cdot \text{CuO} \cdot 2\text{SnO}_2 \cdot 0.78\text{SiO}_2 \cdot 0.27\text{Al}_2\text{O}_3 \cdot n\text{H}_2\text{O}$ ), and dark-green "hydroferstannite" (approx.  $\text{Fe}_2\text{O}_3 \cdot 2.6\text{SnO}_2 \cdot 0.8\text{SiO}_2 \cdot 0.63\text{Al}_2\text{O}_3 \cdot 5\text{H}_2\text{O}$ ). The typical paragenesis of stannite is that with chalcopyrite, cassiterite, topaz, tinnwaldite, and microcline. Often it replaces chalcopyrite and cassiterite, with typical reaction rims, assoc. with older and younger generations of quartz. Chalcopyrite and sphalerite also show regular intergrowths with stannite and bimuthinite indicating unmixing reactions. Late-hydrothermal crystals in stannite are calcite, covellite, and heulandite. The oxidation of stannite was described by W. P. Heiden (1893) and Knipovich (*Zapiski Vsesoiskogo Mineralog. Obshchestva* 75, No. 2 (1940)), with the assumption of a secondary tight-dispersed  $\text{SnO}_2$  or complex Cu-Pb stannate hydrates. The

succession of the oxidation is typically developed: stannite  $\rightarrow$  hydrostannite  $\rightarrow$  hydroferstannite-yellowish "hydroferalite" (about  $0.18\text{Fe}_2\text{O}_3 \cdot 1.78\text{SnO}_2 \cdot \text{Al}_2\text{O}_3 \cdot 3.0\text{H}_2\text{O}$ ). The latter mineral is very similar to Fe halloysite, but somewhat different in its x-ray diagram. All these products are colloidal, very sol., weakly anisotropic, with  $\lambda$  about 1.674 in hydrostannite, 1.752 to 1.774 in hydroferstannite, and 1.705 in hydroferalite. The first step of the oxidation is an enrichment in Sn and  $\text{Fe}^{2+}$ , while Cu and S are removed. The yellow end product contains Al enriched, but no more Sn, Cu, or S; it is evidently only a  $\text{Fe}^{2+}$ -Al silicate-like halloysite, with a slight contamination by scorodite. The authors never observed a change of stannite to  $\text{SnO}_2$ . The rate of the oxidation is about the same for chalcopyrite and stannite. Extensive chem. and spectral-analytical studies showed that all the S in stannite is gradually removed during the oxidation and changed into  $\text{H}_2\text{SO}_4$  and sol. sulfates. Also Sn is entirely removed; Cu is still present in hydrostannite but removed in the later stages of oxidation. Sn is finally removed; Si and Al (derived from feldspars of the mother rock) combine with  $\text{Fe}^{2+}$  to "hydroferalite." W. Eitel.

Dolomanovate.)

sellait from Eastern Transbaikalia. Iv. N. Grigor'ev  
and B. I. Dolomanova. Trudy Mineralog. Muzeyu, Vladivostok.  
Nauk. SSSR, No. 386-41 (1952). —The occurrence is in  
Lower Jurassic clayish schists, aluminites, sandstones, etc.,  
which are intersected by plagiogranite and amznatite-topaz  
quartz veins, with zinnwaldite, fluorite, galena, stannite,  
pyrite, arsenopyrite, etc. In the oxidation zone the usual  
Fe-Mn hydroxides, cerussite, malachite, azurite, and calcite  
are typical. The sellait occurs in the younger tectonic fissures,  
assoc'd. with halloysite, in white crusts; it is uniaxial,  
pos. optical character, cleavage: parallel (100) and (110)  
pool; twins parallel (101), c:c' = 67%; 1.381; n = 1.370;  
ds. 1.44. The chem. analysis and the differential-thermal  
curves show sellait + halloysite in intimate intergrowths;  
the endothermic effects are at 180°, 560°, 900°, and 960°; one  
exothermic effect is at 920°. The paragenesis of sellait with  
halloysite, gennsite (cf. C.R. 49, 138/06), didite, etc.,  
indicates its supogene formation; its Mg content is de-  
rived from the schists which were decompr'd. by V-cong.  
hydrothermal solutions which also brought about the greisen  
formation with topaz and zinnwaldite (cf. V.S. Sobolev,  
Minerolog. Sternik L'vov. Geol. Oshcherni 22, No. 4 (1948)).

W. Eitel

Report from the  
Transbaikalia and its  
metamorphic modification. I.  
Grusley and E. I. Dostoevsky  
Transbaikalia, No. 6, 88-116-  
Tepiz is found mainly in Sn and Au-W deposits in the  
trans-Baikal, in which granites have undergone trans-  
formation into pegmatites. The FISH met. rapids, litho-  
and the nonsolid inclusions are principally gaseous. Met-  
amorphic transformation occurring by vapor found in rock  
metamorphic granites. In granites, under hydrothermal con-  
ditions, results in the introduction of K, Fe, Mg, Ca, Ti,  
and water into the vapor. The inclusions in the modified  
topaz are principally liquid.

C. H. Buchanan

GRIGOR'YEV, Iv.F.; DOLOMANOVA, Ye.I.

Bertrandite from a tin ore deposit in central Transbaikalia.  
Trudy Min.muz. no.7:151-154 '55. (MLRA 9:5)  
(Transbaikalia--Bertrandite)

GRIGOR'YEV, Iv.P.; DOLOMANOVA, Ye.I.

Joseite from a tin ore deposit in central Transbaikalia. Trudy Min.  
muz. no. 7:154-157 '55. (MIRA 9:5)  
(Transbaikalia--Joseite)

✓ Chalcocite-minerals of various types occur in the contacts of carbonaceous rocks and granites. V. P. Kuznetsov, T. N. Gerasimova, L. S. Zhdanov, *Zemle 28*, 24-31 (1953). - Sn-bearing veins occur in sandstone shales, and carbonate rocks near the contacts with biotite and tourmaline biotite granites. Veins in all are the major ore minerals; but whereas quartz, muscovite, topaz, microcline, allanite, and tourmaline are the major gang minerals in veins in the clastic rocks, quartz, fluorite, and tourmaline predominate in the carbonates. Along the granite contacts the carbonates are altered into skarn and skarn-like rocks, and the clastics into hornstone. Late magmatic solns. altered the wall-rocks of the ore veins, the hornstone into greisen, then into mica-schists, talcose and chrysotile rocks, carbonates into fluorite and quartz greisens, and carbonaceous clayey limestones into fluorite-muscovite metamorphics are given.

Wolframite-veins deposited at the contacts of carbonaceous rocks and E. I. Dolgorukov and E. I. Dol-

2  
3  
4  
5  
6  
7  
8  
9  
10  
11  
12  
13  
14  
15  
16  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29  
30  
31  
32  
33  
34  
35  
36  
37  
38  
39  
40  
41  
42  
43  
44  
45  
46  
47  
48  
49  
50  
51  
52  
53  
54  
55  
56  
57  
58  
59  
60  
61  
62  
63  
64  
65  
66  
67  
68  
69  
70  
71  
72  
73  
74  
75  
76  
77  
78  
79  
80  
81  
82  
83  
84  
85  
86  
87  
88  
89  
90  
91  
92  
93  
94  
95  
96  
97  
98  
99  
100  
101  
102  
103  
104  
105  
106  
107  
108  
109  
110  
111  
112  
113  
114  
115  
116  
117  
118  
119  
120  
121  
122  
123  
124  
125  
126  
127  
128  
129  
130  
131  
132  
133  
134  
135  
136  
137  
138  
139  
140  
141  
142  
143  
144  
145  
146  
147  
148  
149  
150  
151  
152  
153  
154  
155  
156  
157  
158  
159  
160  
161  
162  
163  
164  
165  
166  
167  
168  
169  
170  
171  
172  
173  
174  
175  
176  
177  
178  
179  
180  
181  
182  
183  
184  
185  
186  
187  
188  
189  
190  
191  
192  
193  
194  
195  
196  
197  
198  
199  
200  
201  
202  
203  
204  
205  
206  
207  
208  
209  
210  
211  
212  
213  
214  
215  
216  
217  
218  
219  
220  
221  
222  
223  
224  
225  
226  
227  
228  
229  
229  
230  
231  
232  
233  
234  
235  
236  
237  
238  
239  
239  
240  
241  
242  
243  
244  
245  
246  
247  
248  
249  
249  
250  
251  
252  
253  
254  
255  
256  
257  
258  
259  
259  
260  
261  
262  
263  
264  
265  
266  
267  
268  
269  
269  
270  
271  
272  
273  
274  
275  
276  
277  
278  
279  
279  
280  
281  
282  
283  
284  
285  
286  
287  
287  
288  
289  
289  
290  
291  
292  
293  
294  
295  
296  
297  
298  
299  
299  
300  
301  
302  
303  
304  
305  
306  
307  
308  
309  
309  
310  
311  
312  
313  
314  
315  
316  
317  
318  
319  
319  
320  
321  
322  
323  
324  
325  
326  
327  
328  
329  
329  
330  
331  
332  
333  
334  
335  
336  
337  
338  
339  
339  
340  
341  
342  
343  
344  
345  
346  
347  
348  
349  
349  
350  
351  
352  
353  
354  
355  
356  
357  
358  
359  
359  
360  
361  
362  
363  
364  
365  
366  
367  
368  
369  
369  
370  
371  
372  
373  
374  
375  
376  
377  
378  
379  
379  
380  
381  
382  
383  
384  
385  
386  
387  
388  
389  
389  
390  
391  
392  
393  
394  
395  
396  
397  
398  
399  
399  
400  
401  
402  
403  
404  
405  
406  
407  
408  
409  
409  
410  
411  
412  
413  
414  
415  
416  
417  
418  
419  
419  
420  
421  
422  
423  
424  
425  
426  
427  
428  
429  
429  
430  
431  
432  
433  
434  
435  
436  
437  
438  
439  
439  
440  
441  
442  
443  
444  
445  
446  
447  
448  
449  
449  
450  
451  
452  
453  
454  
455  
456  
457  
458  
459  
459  
460  
461  
462  
463  
464  
465  
466  
467  
468  
469  
469  
470  
471  
472  
473  
474  
475  
476  
477  
478  
479  
479  
480  
481  
482  
483  
484  
485  
486  
487  
488  
489  
489  
490  
491  
492  
493  
494  
495  
496  
497  
498  
499  
499  
500  
501  
502  
503  
504  
505  
506  
507  
508  
509  
509  
510  
511  
512  
513  
514  
515  
516  
517  
518  
519  
519  
520  
521  
522  
523  
524  
525  
526  
527  
528  
529  
529  
530  
531  
532  
533  
534  
535  
536  
537  
538  
539  
539  
540  
541  
542  
543  
544  
545  
546  
547  
548  
549  
549  
550  
551  
552  
553  
554  
555  
556  
557  
558  
559  
559  
560  
561  
562  
563  
564  
565  
566  
567  
568  
569  
569  
570  
571  
572  
573  
574  
575  
576  
577  
578  
579  
579  
580  
581  
582  
583  
584  
585  
586  
587  
588  
589  
589  
590  
591  
592  
593  
594  
595  
596  
597  
598  
599  
599  
600  
601  
602  
603  
604  
605  
606  
607  
608  
609  
609  
610  
611  
612  
613  
614  
615  
616  
617  
618  
619  
619  
620  
621  
622  
623  
624  
625  
626  
627  
628  
629  
629  
630  
631  
632  
633  
634  
635  
636  
637  
638  
639  
639  
640  
641  
642  
643  
644  
645  
646  
647  
648  
649  
649  
650  
651  
652  
653  
654  
655  
656  
657  
658  
659  
659  
660  
661  
662  
663  
664  
665  
666  
667  
668  
669  
669  
670  
671  
672  
673  
674  
675  
676  
677  
678  
679  
679  
680  
681  
682  
683  
684  
685  
686  
687  
688  
688  
689  
690  
691  
692  
693  
694  
695  
696  
697  
698  
698  
699  
699  
700  
701  
702  
703  
704  
705  
706  
707  
708  
709  
709  
710  
711  
712  
713  
714  
715  
716  
717  
718  
719  
719  
720  
721  
722  
723  
724  
725  
726  
727  
728  
729  
729  
730  
731  
732  
733  
734  
735  
736  
737  
738  
739  
739  
740  
741  
742  
743  
744  
745  
746  
747  
748  
749  
749  
750  
751  
752  
753  
754  
755  
756  
757  
758  
759  
759  
760  
761  
762  
763  
764  
765  
766  
767  
768  
769  
769  
770  
771  
772  
773  
774  
775  
776  
777  
778  
779  
779  
780  
781  
782  
783  
784  
785  
786  
787  
788  
788  
789  
790  
791  
792  
793  
794  
795  
796  
797  
798  
798  
799  
799  
800  
801  
802  
803  
804  
805  
806  
807  
808  
809  
809  
810  
811  
812  
813  
814  
815  
816  
817  
818  
819  
819  
820  
821  
822  
823  
824  
825  
826  
827  
828  
829  
829  
830  
831  
832  
833  
834  
835  
836  
837  
838  
839  
839  
840  
841  
842  
843  
844  
845  
846  
847  
848  
849  
849  
850  
851  
852  
853  
854  
855  
856  
857  
858  
859  
859  
860  
861  
862  
863  
864  
865  
866  
867  
868  
869  
869  
870  
871  
872  
873  
874  
875  
876  
877  
878  
879  
879  
880  
881  
882  
883  
884  
885  
886  
887  
888  
888  
889  
889  
890  
891  
892  
893  
894  
895  
896  
897  
898  
898  
899  
899  
900  
901  
902  
903  
904  
905  
906  
907  
908  
909  
909  
910  
911  
912  
913  
914  
915  
916  
917  
918  
919  
919  
920  
921  
922  
923  
924  
925  
926  
927  
928  
929  
929  
930  
931  
932  
933  
934  
935  
936  
937  
938  
939  
939  
940  
941  
942  
943  
944  
945  
946  
947  
948  
949  
949  
950  
951  
952  
953  
954  
955  
956  
957  
958  
959  
959  
960  
961  
962  
963  
964  
965  
966  
967  
968  
969  
969  
970  
971  
972  
973  
974  
975  
976  
977  
978  
979  
979  
980  
981  
982  
983  
984  
985  
986  
987  
988  
988  
989  
989  
990  
991  
992  
993  
994  
995  
996  
997  
997  
998  
998  
999  
999  
1000  
1001  
1002  
1003  
1004  
1005  
1006  
1007  
1008  
1009  
1009  
1010  
1011  
1012  
1013  
1014  
1015  
1016  
1017  
1018  
1019  
1019  
1020  
1021  
1022  
1023  
1024  
1025  
1026  
1027  
1028  
1029  
1029  
1030  
1031  
1032  
1033  
1034  
1035  
1036  
1037  
1038  
1039  
1039  
1040  
1041  
1042  
1043  
1044  
1045  
1046  
1047  
1048  
1049  
1049  
1050  
1051  
1052  
1053  
1054  
1055  
1056  
1057  
1058  
1059  
1059  
1060  
1061  
1062  
1063  
1064  
1065  
1066  
1067  
1068  
1069  
1069  
1070  
1071  
1072  
1073  
1074  
1075  
1076  
1077  
1078  
1079  
1079  
1080  
1081  
1082  
1083  
1084  
1085  
1086  
1087  
1088  
1088  
1089  
1089  
1090  
1091  
1092  
1093  
1094  
1095  
1096  
1097  
1098  
1098  
1099  
1099  
1100  
1101  
1102  
1103  
1104  
1105  
1106  
1107  
1108  
1109  
1109  
1110  
1111  
1112  
1113  
1114  
1115  
1116  
1117  
1118  
1119  
1119  
1120  
1121  
1122  
1123  
1124  
1125  
1126  
1127  
1128  
1129  
1129  
1130  
1131  
1132  
1133  
1134  
1135  
1136  
1137  
1138  
1139  
1139  
1140  
1141  
1142  
1143  
1144  
1145  
1146  
1147  
1148  
1149  
1149  
1150  
1151  
1152  
1153  
1154  
1155  
1156  
1157  
1158  
1159  
1159  
1160  
1161  
1162  
1163  
1164  
1165  
1166  
1167  
1168  
1169  
1169  
1170  
1171  
1172  
1173  
1174  
1175  
1176  
1177  
1178  
1179  
1179  
1180  
1181  
1182  
1183  
1184  
1185  
1186  
1187  
1188  
1188  
1189  
1189  
1190  
1191  
1192  
1193  
1194  
1195  
1196  
1197  
1198  
1198  
1199  
1199  
1200  
1201  
1202  
1203  
1204  
1205  
1206  
1207  
1208  
1209  
1209  
1210  
1211  
1212  
1213  
1214  
1215  
1216  
1217  
1218  
1219  
1219  
1220  
1221  
1222  
1223  
1224  
1225  
1226  
1227  
1228  
1229  
1229  
1230  
1231  
1232  
1233  
1234  
1235  
1236  
1237  
1238  
1239  
1239  
1240  
1241  
1242  
1243  
1244  
1245  
1246  
1247  
1248  
1249  
1249  
1250  
1251  
1252  
1253  
1254  
1255  
1256  
1257  
1258  
1259  
1259  
1260  
1261  
1262  
1263  
1264  
1265  
1266  
1267  
1268  
1269  
1269  
1270  
1271  
1272  
1273  
1274  
1275  
1276  
1277  
1278  
1279  
1279  
1280  
1281  
1282  
1283  
1284  
1285  
1286  
1287  
1288  
1288  
1289  
1289  
1290  
1291  
1292  
1293  
1294  
1295  
1296  
1297  
1298  
1298  
1299  
1299  
1300  
1301  
1302  
1303  
1304  
1305  
1306  
1307  
1308  
1309  
1309  
1310  
1311  
1312  
1313  
1314  
1315  
1316  
1317  
1318  
1319  
1319  
1320  
1321  
1322  
1323  
1324  
1325  
1326  
1327  
1328  
1329  
1329  
1330  
1331  
1332  
1333  
1334  
1335  
1336  
1337  
1338  
1339  
1339  
1340  
1341  
1342  
1343  
1344  
1345  
1346  
1347  
1348  
1349  
1349  
1350  
1351  
1352  
1353  
1354  
1355  
1356  
1357  
1358  
1359  
1359  
1360  
1361  
1362  
1363  
1364  
1365  
1366  
1367  
1368  
1369  
1369  
1370  
1371  
1372  
1373  
1374  
1375  
1376  
1377  
1378  
1379  
1379  
1380  
1381  
1382  
1383  
1384  
1385  
1386  
1387  
1388  
1388  
1389  
1389  
1390  
1391  
1392  
1393  
1394  
1395  
1396  
1397  
1398  
1398  
1399  
1399  
1400  
1401  
1402  
1403  
1404  
1405  
1406  
1407  
1408  
1409  
1409  
1410  
1411  
1412  
1413  
1414  
1415  
1416  
1417  
1418  
1419  
1419  
1420  
1421  
1422  
1423  
1424  
1425  
1426  
1427  
1428  
1429  
1429  
1430  
1431  
1432  
1433  
1434  
1435  
1436  
1437  
1438  
1439  
1439  
1440  
1441  
1442  
1443  
1444  
1445  
1446  
1447  
1448  
1449  
1449  
1450  
1451  
1452  
1453  
1454  
1455  
1456  
1457  
1458  
1459  
1459  
1460  
1461  
1462  
1463  
1464  
1465  
1466  
1467  
1468  
1469  
1469  
1470  
1471  
1472  
1473  
1474  
1475  
1476  
1477  
1478  
1479  
1479  
1480  
1481  
1482  
1483  
1484  
1485  
1486  
1487  
1488  
1488  
1489  
1489  
1490  
1491  
1492  
1493  
1494  
1495  
1496  
1497  
1498  
1498  
1499  
1499  
1500  
1501  
1502  
1503  
1504  
1505  
1506  
1507  
1508  
1509  
1509  
1510  
1511  
1512  
1513  
1514  
1515  
1516  
1517  
1518  
1519  
1519  
1520  
1521  
1522  
1523  
1524  
1525  
1526  
1527  
1528  
1529  
1529  
1530  
1531  
1532  
1533  
1534  
1535  
1536  
1537  
1538  
1539  
1539  
1540  
1541  
1542  
1543  
1544  
1545  
1546  
1547  
1548  
1549  
1549  
1550  
1551  
1552  
1553  
1554  
1555  
1556  
1557  
1558  
1559  
1559  
1560  
1561  
1562  
1563  
1564  
1565  
1566  
1567  
1568  
1569  
1569  
1570  
1571  
1572  
1573  
1574  
1575  
1576  
1577  
1578  
1579  
1579  
1580  
1581  
1582  
1583  
1584  
1585  
1586  
1587  
1588  
1588  
1589  
1589  
1590  
1591  
1592  
1593  
1594  
1595  
1596  
1597  
1598  
1598  
1599  
1599  
1600  
1601  
1602  
1603  
1604  
1605  
1606  
1607  
1608  
1609  
1609  
1610  
1611  
1612  
1613  
1614  
1615  
1616  
1617  
1618  
1619  
1619  
1620  
1621  
1622  
1623  
1624  
1625  
1626  
1627  
1628  
1629  
1629  
1630  
1631  
1632  
1633  
1634  
1635  
1636  
1637  
1638  
1639  
1639  
1640  
1641  
1642  
1643  
1644  
1645  
1646  
1647  
1648  
1649  
1649  
1650  
1651  
1652  
1653  
1654  
1655  
1656  
1657  
1658  
1659  
1659  
1660  
1661  
1662  
1663  
1664  
1665  
1666  
1667  
1668  
1669  
1669  
1670  
1671  
1672  
1673  
1674  
1675  
1676  
1677  
1678  
1679  
1679  
1680  
1681  
1682  
1683  
1684  
1685  
1686  
1687  
1688  
1688  
1689  
1689  
1690  
1691  
1692  
1693  
1694  
1695  
1696  
1697  
1698  
1698  
1699  
1699  
1700  
1701  
1702  
1703  
1704  
1705  
1706  
1707  
1708  
1709  
1709  
1710  
1711  
1712  
1713  
1714  
1715  
1716  
1717  
1718  
1719  
1719  
1720  
1721  
1722  
1723  
1724  
1725  
1726  
1727  
1728  
1729  
1729  
1730  
1731  
1732  
1733  
1734  
1735  
1736  
1737  
1738  
1739  
1739  
1740  
1741  
1742  
1743  
1744  
1745  
1746  
1747  
1748  
1749  
1749  
1750  
1751  
1752  
1753  
1754  
1755  
1756  
1757  
1758  
1759  
1759  
1760  
1761  
1762  
1763  
1764  
1765  
1766  
1767  
1768  
1769  
1769  
1770  
1771  
1772  
1773  
1774  
1775  
1776  
1777  
1778  
1779  
1779  
1780  
1781  
1782  
1783  
1784  
1785  
1786  
1787  
1788  
1788  
1789  
1789  
1790  
1791  
1792  
1793  
1794  
1795  
1796  
1797  
1798  
1798  
1799  
1799  
1800  
1801  
1802  
1803  
1804  
1805  
1806  
1807  
1808  
1809  
1809  
1810  
1811  
1812  
1813  
1814  
1815  
1816  
1817  
1818  
1819  
1819  
1820  
1821  
1822  
1823  
1824  
1825  
1826  
1827  
1828  
1829  
1829  
1830  
1831  
183

Dolomanova, I. T.

GRIGOR'YEV, Iv. P.; DOLOMANOVA, I. T.

Genetic types of tin ore deposits in Transbaikalia. Trudy MGRI 29:  
38-51 "56.  
(Transbaikalia--Tin ores)

(MIRA 10:4)

DOLOMANOVA, YE. I.

15-1957-7-9275

Translation from: Referativnyy zhurnal, Geologiya, 1957, Nr 7,  
p 71 (USSR)

AUTHOR: Dolomanova, Ye. I.

TITLE: Assimilation and Granitization of Sandstone-Shale Rocks  
by Hypabyssal Granitic Intrusions (K voprosu ob assimil-  
yatsii i granitizatsii peschaniko-slantsevykh porod  
granitnymi intruziyami gipabissal'nogo tipa).

PERIODICAL: Sov. geologiya, vol 51, 1956, pp 113-141

ABSTRACT: An intrusive granite porphyry situated between the right-  
hand tributaries of the Ingoda River--the Sluchaynyy and  
Listvenichnyy Springs--is younger than the Paleozoic  
sedimentary sandstone-shale rocks and the great granite  
intrusion of old Kimmeriyskiy age; it was intruded along  
regional zones of deformation of folds and faults. The  
intrusive was formed in a complex tectonic environment  
from magma rich in volatile constituents, particularly  
in B and F; these volatiles produced widespread assimila-

Card 1/3

15-1957-7-9275

Assimilation and Granitization of Sandstone-Shale Rocks by Hypabyssal  
Granitic Intrusions (Cont.)

lation, contamination, and granitization in the endogene and exogene contact zones of the intrusion. The rocks of the main intrusive phase are chiefly porphyries with characteristic networks of veins of quartz and microcline-perthite segregations which were produced by earlier autopneumatolysis of the granite porphyries. Biotite granites, leucocratic granites, and fine-grained biotite granites also belong to the main intrusive phase. Contaminated granitoid rocks have developed chiefly in the endogene zone of the intrusion, characterized by variable structure, texture, and mineral composition, especially along gently inclined contacts. Sandstone-shale xenoliths are distributed irregularly in the mass and display various degrees of assimilation and recrystallization. Granitization of hornfelses is manifested chiefly in the formation of quartz and microcline-perthite porphyroblasts, in the appearance of topaz and fluorite, and in the recrystallization of biotite. Reaction rims of albite, quartz, biotite, and fluorite (sometimes of apatite), or of ilmenite or coarsely platy biotite, form about the hornfels

Card 2/3

Assimilation and Granitization of Sandstone-Shale Rocks by Hypabyssal  
Granitic Intrusions (Cont.) 15-1957-7-9275

fragments. The hybrid granites have been enriched in biotite; the plagioclase is more abundant and more basic; and the content of V, Ni, Cr, and Co has increased. The most strongly contaminated rocks show both granulitic and poikilitic textures. Mafic porphyries; these contain, in association with quartz, albite, topaz, fluorite, and tourmaline, the ore minerals wolframite, molybdenite, and arsenopyrite, which formed by interaction between the granitic magma and the hornfels xenoliths. It is shown that there is a high content of Sn in the autopneumatolitically altered "normal" granites, a low content in the hybrid rocks, and a decrease in content in the host rocks proportional to their distance away from the contact with the intrusion.

Card 3/3

O. V. Bryzgalin

Translation from: Referativnyy zhurnal, Geologiya, 1957, Nr 5,  
p 112 (USSR) 15-57-5-6499

AUTHORS: Grigor'yev, Iv. F., Dolomanova, Ye. I.

TITLE: The Age and Structural Position of the Granite  
Intrusions and the Tin-Tungsten Deposits Associated  
With Them in the Central Trans-Baikal Region (O  
vozraste i strukturnom polozenii granitnykh intruziy  
i svyazannykh s nimi olovyanno-volframovykh mestoro-  
zhdeniy v Tsentral'nom Zabaykal'ye)

PERIODICAL: Tr. In-ta geol. rud. mestorozhd. petrogr. mineralogii  
i geokhimii, 1956, Nr 3, pp 236-269

ABSTRACT: The following intrusive formations are found in the  
central trans-Baikal region [the Zachikinskaya tayga  
(taiga) and the Daurskiy khrebet (Range)]; 1) Variscan  
granodiorites and ultrabasic rocks, not widespread;  
2) early Cimmerian (early Mesozoic) biotite granites

Card 1/4

15-57-5-6499  
The Age and Structural Position of the Granite (Cont.)

and other varieties forming the "Great Intrusion" (60 percent of the area of the central trans-Baikal region), dated by monazite and samarskite as 180 to 190 million years old (Early Triassic); and 3) late Cimmerian granites and granite porphyries, represented by small intrusions. These last intrusions contain fragments of partly assimilated vein quartz and kersantite as well as fragments of biotite granite of early Cimmerian age. The geological and structural position of the granitoidal intrusions of early and late Cimmerian age, which has a direct relationship with the tin-tungsten mineralization, associated with greisenization, occurs in those zones where these intrusions form cupola-like extensions into the country rock. A study of the jointing in the Shumilovo region in the "Great Intrusion" has shown that the aggregate of gently dipping fractures represents a series of gently inclined waves proceeding in two mutually perpendicular directions and forming at their intersections a system of dome-like bends. The greisen and ore deposits are confined to these zones. The late Cimmerian granitoidal intrusions

Card 2/4

15-57-5-6499

The Age and Structural Position of the Granite (Cont.)

show control by large north-south and northeasterly fractures superimposed on Paleozoic rocks and early Cimmerian intrusions. This is demonstrated by a study of the jointing in the region of the Ingoda fracture (a fault) where it is seen that the system of joints associated with the fracture are similar in the Paleozoic sandy shales and in the granitoidal rocks of the "Great Intrusion" but not present in the granite porphyries. The tin-tungsten mineralization of the region is of two ages: it is genetically associated both with the early Cimmerian intrusive complex and with the late Cimmerian granitoidal masses. The older mineralization is localized in the cupola-like extensions of the intrusions, but the younger mineralization is associated with feather-type joints associated with regional fractures. The deposits generally occur in the country rock over protuberances of the intrusives. Ore-bearing granitoidal rocks of both ages are over-saturated in silica and alumina, under-saturated in lime, and are distinguished by a high content of fluorine. These distinguishing features are especially strong in Card 3/4

The Age and Structural Position of the Granite (Cont.) 15-57-5-6499

the late Cimmerian granite porphyries. The authors state their opinion that the early and late Cimmerian granitoidal rocks are genetically related and are derived from the same magmatic source.

Card 4/4

Ye. P. M.

15-57-4-4899

Translation from: Referativnyy zhurnal, Geologiya, 1957, Nr 4,  
p 123 (USSR)

AUTHORS: Grigor'yev, Iv. F., Dolomanova, Ye. I.

TITLE: Tin Ores Transitional Between Cassiterite-Quartz and  
Cassiterite-Sulfide (Ob olovorudnykh mestorozhdeniyakh  
perekhodnykh tipov mezhdu mestorozhdeniyami kassiter-  
ito-kvartsevoy i kassiterito-sul'fidnoy formatsii)

PERIODICAL: Tr. Kn-ta geol. rud. mestorozhd. petrogr., mineralogii  
i geokhimii, 1956, Nr 3, pp 279-301

ABSTRACT: The authors have arrived at the conclusion that tin  
ores which are transitional between cassiterite-  
quartz and cassiterite-sulfide formations should be  
classed as a separate cassiterite-quartz-sulfide for-  
mation. These formations are different in a number  
of geological and mineralogical characteristics. The  
ore-bearing intrusives of granitoids, with which such  
deposits are genetically associated, are intruded

Card 1/3

15-57-4-4899

Tin Ores Transitional (Cont.)

along large regional tectonic dislocations; the fissures produced by these dislocations contain the ore. The formations are associated with the zone of exocontact and lie in sandstone-shale rocks, granites, effusives, etc. The ore-bearing intrusives are of small dimensions and are of the "fissure" type. In composition they represent granite-porphyrries, or granodiorite-porphyrries (the Little Khingan Mountains), and granodiorites (Yana-Adychi region). At the same time, a direct genetic connection of the cassiterite-quartz-sulfide formations with the intrusives has been established only for certain locations in the trans-Baikal area; here the association is with granite-porphyrries. Tin ore deposits of the cassiterite-quartz-sulfide formation are represented by network zones, stockwork zones, brecciated zones, fissured zones, and lenses. The mineral composition of the ores in the cassiterite-quartz-sulfide formations is unique, since it includes minerals characteristic of both the cassiterite-quartz and the cassiterite-sulfide formations. Chief minerals are:

Card 2/3

Tin Ores Transitional (Cont.)

15-57-4-4899

1) ore minerals -- arsenopyrite, pyrrhotine, sphalerite, galena, chalcopyrite, cassiterite, tungstenite, and scheelite; 2) non-ore minerals -- tourmaline, chlorite, muscovite, quartz, topaz, and fluorite. Ores of the given formation are multi-stage. Changes in the vicinity of the veins are expressed in greisenization, tourmalinization, chloritization, sericitization, and quartzification. Thus the ores in these formations are characterized by lack of the genetic features of the cassiterite-quartz and cassiterite-sulfide formations. This is caused not only by the composition of the postmagmatic solutions themselves, but also by the geologic and structural environment in which these formations originated. Industrially, tin ores of the transitional type represent large reserves of low-grade ore; they sometimes contain tungstenite and other values.

Card 3/3

Ye. P. M.

GRIGOR'YEV, Iv.P.; DOLOMANOVA, Ye.I.;

Relation of tin mineralization to dikes in Transbaikalia. Izv.AN  
SSSR, Ser.geol. 22 no.1:67-89 Ja '57. (MLRA 10:3)

1. Institut geologii rudnykh mestoroshdeniy, petrografii, mineralogii  
i geokhimii AN SSSR, Moskva.  
(Transbaikalia--Tin ores) (Dikes (Geology))

Geological setting of the Saimovskite localities  
belong to group 1 (Ural, Orenburg).  
**Zotik-Pazyryk, Ural, Orenburg.** In western Trans-Uralia, the Saimovskite deposit of Sotykhovka is characterized by the occurrence of cassiterite-topaz-quartz-ammonite veins; a peculiar new mineral was observed, called sainovskite. The country rocks are metamorphic Lower Jurassic: jaspe, gneiss, and vein veins in contact with amphibole granites, plagioperthites, and orthopyroxene porphyries. The veins in the veins contain cassiterite, wolframite, columbite, with zinnwaldite, followed by rutile, apatite, magnetite, fluorite, staurolite, pyrochlore, perovskite, chalcocite, pyrite, sphalerite, galena, and calcite. In the oxidation zone are kaolinite, halloysite, dickite, iron-muscovite schist, geckosilite, celadonite, arfvedsonite, rhoibrookite, lepidolite, and zeunerite. Sainovskite occurs in two generations, the first (high) generation on sills and in a typical pegmatitic high-temp. pegmatites with cassiterite, topaz, and quartz, also crystals between scaly zinnwaldite. Characteristic is the replacement of thin smirnovskite by topaz and cassiterite, or secondary minerals. The second, low-temp., generation shows sainovskite crystals around cassiterite, or intergrown with it, or inclusions in fibrous galena, and triplite. Sainovskite shows indistinct, perhaps tetragonal, crystal form of prismatic habit, usually in aggregate, not larger than 1 X 1 to 3 X 5 mm. Color is variable with different states of weathering, from dark brown to red-orange, or yellow-brown, and even nearly colorless. Often the effect is banding and striated. Sainovskite is a typical metamorphic product of an unknown mineral, highly hydrazidic, fatty luster, fracture conchoidal, no cleavage, hardness about 4; d. 4.03; isotropic, with occasional anisotropic portions, with  $\gamma = 1.882$ ;  $\alpha = 1.878$ ; very low birefringence, uniaxial, pos. In the dark brown crystals  $\alpha$  is about 1.70-1.71/18. The differential thermal curves show dehydration effects between 100° and 250°, all the H<sub>2</sub>O is lost at 50°; between 300° and

1100° also distinct loss of F and a slight increase are observed. A slight smoothening effect between 60° and 70° may indicate the recrystallization of the metanite substance (X-ray amorphous; above 110° distinct lines of thoriteite ( $\text{Ca}_2\text{Si}_2\text{O}_5$ ) appear indicating an intergrowth of compns. Four chem. analyses are given, which show the highly complex compn. of Smilnevalite. It contains rare earths; the analyses correspond to the type formulae ( $\text{Ca}, \text{Ce}, \text{La}, (\text{OH})_2(\text{Fe}^{2+}, \text{Mn}^{2+})\text{Si}_2\text{O}_5(\text{OH})_2$ ), or the same with 0.3 mols. H<sub>2</sub>O. The metanite group includes the complex phosphato-aluminosilicate type; F<sup>-</sup> and OH<sup>-</sup> are also variable. In the cations, the replacement of Th by Ca, further by Ce, La, Cd... is also characteristic. The mineral is easily changed to orange-yellow hydrous silicates (it may decompose). This development is illustrated by the results of sequential, successive, and spectral analysis of the weathering residue. X-ray spectrograph: leucite-like enrichment in Li, Pb, Nd, and U; (a) the colorless spinelike variety. This shows a strong green fluorescence in ultraviolet light whereas the dark brown mineral does not show any luminescence (exceptional effect of Fe content). Smilnevalite is composed with thorite but it contains less ThO<sub>3</sub> but more rare earths and F<sup>-</sup> than this. - The H<sub>2</sub>O contents are similar in both; smilnevalite, however, contains much F<sup>-</sup> which is absent in thorite, and this contains CO<sub>3</sub> which is absent in smilnevalite. In its exterior aspect, another so-called smilnevalite from pegmatites of eastern Khibabian resembles metathorite. In Nb one like ambergisite, or orange (thorite) from Norway. This smilnevalite variety has no rare earths, and also no Cs<sub>2</sub>, but contains U.

4  
14E 4/

DOLGANOVA, Yelizaveta Ivanovna; SHIPULIN, F.K., otd. red.; IVANOV, B.V.,  
red.ind.-va; SHEVCHENKO, G.M., tekhn.red.

[Ingoia tin-wolframium deposits and their genetic characteristics]  
Olovianno-vol'framovye mestorozhdeniya Ingodinskogo rudnogo uzla i  
ikh geneticheskie osobennosti. Moskva, Izd.-vo Akad.nauk SSSR, 1959.  
297 p. (Akademika nauk SSSR. Institut geologii rudnykh mestorozhdenii,  
petrografii, mineralogii i geokhimii. Trudy, no.23) (MIRA 13:2)  
(Ingoda Valley--Tin ores)  
(Ingoda Valley--Wolframite ores)

GRIGOR'YEV, I.F.; DOLOMANOVA, Ye.I.

Gearksite. Trudy Min.muz. no.10:185-186 '59.  
(Gearksite) (Transbaikalia---Gearksutite) (MIRA 16:8)

S/015/60/000/009/004/005  
A052/A129

AUTHOR: Dolomanova, Ye. I.

TITLE: Tin-tungsten deposits of the Ingodinsk mining field and their genetic characteristics

PERIODICAL: Referativnyy zhurnal Geologiya, 1960, no. 9, 299, abstract 17792  
(Tr. In-ta geol. rudn. mestorozh. petrogr., mineralogii i geo-khimii. AN SSSR. 1959, no. 23, 299 pages, illustrations, maps).

TEXT: Tin-tungsten deposits of the Ingodinsk mining field are located in the rare metal zone of the Transbaikal region. A short geological and petrographical description of the area and its tectonic structure is given; the petrochemical characteristics of the tin-bearing granitoids and of the sand-schist rocks of the meso-upperpaleozoic era containing them are investigated; the Ingodinsk tin deposits, the Levo-Ingodinsk tin-tungsten deposits and the "Novoye" molybdenum-tin-tungsten deposits are described. The genesis of the Ingodinsk deposits and their place in the general genetic classification of tin deposits are analyzed. The deposits of the Ingodinsk mining field are connected genetically with the neokimeridgian granite-porphyrles. This connection is proved by the

Card 1/2

YAKZHIN, Aleksandr Andreyevich, doktor geol-min.nauk, prof.(1907-1961);  
DOLOMANOVA, Ye.I., red.; OVCHINNIKOV, S.V., red. izd-va;  
BYKOVA, V.V., tekhn. red.

[Characteristics of the distribution and formation of fluorite  
deposits in Transbaikalia] Zakonomernosti razmeshcheniya i for-  
mirovaniya fluoritovykh mestorozhdenii Zabaikal'ia. Moskva,  
Gosgeoltexhizdat, 1962. 249 p. (MIRA 15:9)  
(Transbaikalia--Fluorite)

DOLONANOVA, Ye.I.

Zeolites from the Shirlovaya Gora deposit in eastern Transbaikalia.  
Trudy Min.muz. no.13:172-178 '62. (MIRA 16:2)  
(Transbaikalia--Zeolites)

DOLOMANOVA, Ye.I.; NESTEROVA, Yu.B.; ARAPOVA, G.A.

Tl and Sn containing boudantite from the Bol'shaya Shirlowaya  
Gora deposit (eastern Transbaikalia). Trudy Min.muz. no.13:179--  
190 '62. (MIRA 16:2)  
(Transbaikalia—Boudantite)

DOLOMANOVA, Ye.I.; MOLEVA, V.A.

Danalite from the deposit of the cassiterite-quartz-sulfide  
formation in Siberia. Trudy Min. muz. no.14:215-218 '63.  
(MIRA 16:10)  
(Siberia--Danalite)

DOLOMANOVA, Ye.I.

Lead-zinc mineralization in some cassiterite-quartz-sulfide  
deposits of eastern Transbaikalia. Trudy IGEM no. 83:468-504  
'63. (MIRA 16:11)

ZVYAGIN, B.B.; DOLMANOVA, Ye.I.; SOBOLEVNA, S.V.; MOLEVA, V.A.

Dioctahedral Al<sub>2</sub>Sn<sub>3</sub>O<sub>10</sub> from the Lovo-Ingedinsk tin-tungsten deposit in Transbaikalia. Dokl. AN SSSR 165 no. 2:410-412 N 1965.  
(MIRA 18:11)

1. Submitted May 21, 1965.

SOLOMONOV, YE. V.

"Hygienic characteristics of the organization of open-air days in  
creches and kindergartens."

report submitted at the 13th All-Union Congress of Hygienists, Epidemiologists  
and Infectionists, 1959.

DOLOMINO, N., red.; YISENKO, A., tekhn.red.

[Put chemistry to work for the good of the Soviet people;  
a collection of papers] Khimiia - na blago sovetskogo  
cheloveka; sbornik materialov. Simferopol', Krymizdat, 1958.  
72 p. (MIRA 12:4)  
(Chemical industries)

BLUDILIN, Mikhail Aver'yanchich; DOLOMINO, N., red.; FISENKO, A.,  
telhn. red.

[Main potential] Glavnyi rezerv. Simferopol', Krymizdat, 1960.  
15 p. (MIRA 15:11)

1. Glavnyy inzhener Kamyshturinskogo zhelezorudnogo kombinata  
(for Bludilin).  
(Kerch Peninsula--Iron and steel plants---Technological in-  
novations)

ERYANOV, Vasiliy Vladimirovich; DOLOMINO, N., red.; FISENKO, A.,  
tekhn. red.

[Extracting large blocks] Dobycha krupnykh blokov. Simferopol',  
Krymizdat, 1961. 44 p.  
(MIRA 15:11)  
(Crimea--Building stones)

KHOMENKO, Ivan Andreyevich; DOLOMINO, N., red.; POLYAKOV, I., red.;  
ISUPOVA, N., tekhn. red.

[Making every minute count] Droozha kazhdoi minutoi. Simferopol',  
Krymizdat, 1961. 29 p. (MIRA 15:12)

1. Direktor Simferopol'skogo kozhevenno-obuvnogo kombinata im.  
F.Ye.Dzerzhinskogo (for Khomenko).  
(Simferopol'--Shoe industry--Labor productivity)

FESHCHENKO, Petr Stepanovich, kand. ekon. nauk; DOLOMINO, N., red.;  
POLYAKOV, I., red.; FISENKO, A., tekhn. red.

[Towards a new upsurge in state farm economy] K novomu pod"  
emu ekonomiki sovkhozov; v ponoshch' izuchaiushchim ekono-  
miku sel'skogo khoziaistva. Simferopol', Krymizdat, 1961. 54 p.  
(MIRA 15:11)

(Crimea—State Farms)

SOBCHUK, B.A.; DOLOSHITSKIY, L.M. [Doloshyts'kyi, L.M.]; GLADYSHEVSKAYA,  
T.N. [Hladyshevs'ka, T.M.]

Carboxymyoglobin in rats during acute carbon monoxide poisoning.  
Ukr. biokhim. zhur. 33 no.6:848-855 '61. (MIRA 14:12)

J. Department of Biochemistry and Department of General Hygiene  
of Lvov Medical Institute.  
(CARBON MONOXIDE PHYSIOLOGICAL EFFECT) (MYOGLOBIN)

VAL'TUKH, E.; DOLOTENKOVA, L.

Fundamental advantages of socialism in utilizing production  
funds. Fin. SSSR 22 no.3:48-57 Mr '61. (MIRA 14:7)  
(Capital) (Communism)

ABALKIN, Leonid Ivanovich; VAL'TUKH, Konstantin Kurtovich;  
DOLOFENKOVA, Liliya Pavlovna; MANDRYGINA, Faina  
Aleksandrovna; PLYSHEVSKIY, B.P., red.; MATSUK, R.V.,  
red. izd-va; GARINA, T.D., tekhn. red.

[Study of the production of the means of production under the  
conditions of the general crisis of capitalism; based on the  
U.S.A.] Ocherk vosprievodstva v usloviyah obshchego krisisa  
kapitalizma; na primere SShA [By] L.I. Abalkin i dr. Moskva,  
Vysshiaia shkola, 1962. 118 p. (MIRA 15:3)  
(United States—Economic conditions)

DOLOTIKHINA, Z. G.

Dolotikhina, Z. G. -- "The Interrelationship between High Rye Grass and Its Components When It is Introduced into the Field Grass Mixtures in Belgorod Oblast." Leningrad Order of Lenin State University A. A. Zhdanov. Leningrad, 1956. (Dissertation For the Degree of Candidate in Biological Sciences).

So. Knizhnaya Letopis', No. 11, 1956, pp 103-114

KHODZHAYEV, Aqzam Ashrapovich; DOLONIN, V., red.; MEL'NIKOV, A., tekred.

[Food industry of Uzbekistan under the seven-year plan] Pishchevaya promyshlennost' Uzbekistana v semiletke. Tashkent, Gos. izd-vo Uzbekskoi SSR, 1959. 61 p. (MIMA 13:8)  
(Uzbekistan--Food industry)

DOLOTKAZIN, A.A., fel'dsher (Dagomys Krasnodarskogo kraya)

Case of mutism caused by sulphur plugs in the ears. Fel'd. 1  
akush. 26 no. 3:47 Mr '61. (MIRA 14:3)  
(CHILDREN, DEAF) (EAR—CARE AND HYGIENE)

USSR/Cultivated Plants. Technical Plants. Oil and II  
Sugar Bearing Plants.

Abs Jour : Ref Zhur-Biol., No 15, 1958, 63263

Author : Dalotkazin, I.

Inst. : Kara-Kum Experiment and Amelioration Station  
of the All-Union Scientific Research Institute  
of Hydraulic Engineering and Amelioration.

Title : An Experiment in Sprinkling Fine-Fiber Cotton  
in Turkmenistan.

Orig Pub : Khlopkovodstvo, 1957, No 5, 51-54

Abstract : In 1956, at the Moskva kolkhoz in Mary Oblast,  
a field experiment was conducted by the Kara-  
Kum Experiment and Amelioration Station of the  
All-Union Scientific Research Institute of Hy-  
draulic Engineering and Amelioration. Sprin-  
kling increased cotton yields by 3.7 centners/

Card : 1/2

USSR/Cultivated Plants. Technical Plants. Oil and M  
Sugar Bearing Plants.

Abs Jour : Ref Zhur-Biol., № 15, 1958, 68263

hectare, as compared with natural irrigation,  
and the expenditure of water was significantly  
lower. Plants developed similarly under both  
methods of irrigation, but on the sprinkled  
plots the plants preserved a larger number  
of seedcases. -- A. M. Smirnov

Card : 2/2

112

DOLOTKAZIN, I.Ya., insh.

Results of experiments in sprinkler irrigation of cotton. Trudy  
VNIIGIM 32:50-60 '59. (MIRA 13:8)  
(Cotton--Irrigation) (Sprinkler irrigation)

18

New technology of building gray 1000 ft. long Preformed  
Building and 1000 ft. long Preformed Building  
1989. No. 8,200-1,000 cubic meters were equipped with a  
water-cooled steel and turbine air compressors. This per-  
mitted increasing their production with 10-15% during op-  
eration. Details are given both of the equipment and opera-  
tion.

J. E. G.

AUTHOR:

Dolotov, G.P.  
Zhuravlev, P.A.  
Kuznetsov, I.I  
Kogan, G.M.  
Kondakov, Ye.A.  
Nesterenko, P.S.

SOV/94-58-11-9/28

TITLE:

The Installation of a Radiation Recuperator on a Cupola  
(Ustanovka radiatsionnogo rekuperatora na vayranke)

PERIODICAL: Promyshlennaya Energetika, 1958, Nr 11, p 19. (USSR)

ABSTRACT: This suggestion was awarded a fifth premium in an All-Union Power Economy competition. Hitherto little use has been made of waste heat from foundry cupolas largely because the heat exchangers become dirty very quickly and therefore inefficient. Metal radiation recuperators of simple construction have recently been used abroad for this purpose. The authors proposed the installation of radiation recuperators for heating blast air on two cupolas of 18 tons per hour upwards. A sketch of the equipment is given. The recuperator consists of two metal tubes with an annular gap of

Card 1/2

SOV/94-58-11-9/23

The Installation of a Radiation Recuperator on a Cupola

32 mm; the recuperator is 6,000 mm high and constructional details are given. The method of installing the device is briefly described. The equipment has proved satisfactory in service and economises about 1,180 tons of coke a year. There is 1 figure.

Card 2/2

*Dekotov, G.F.*

22(5) FILED 1960 SEPTEMBER 2007 BY THE GOVERNMENT OF THE SOVIET UNION  
FOR RELEASE

Central Scientific Research Institute of Machine-Building  
Industry of Industrial Plants, Ministry of Machine-Building Industry  
of Soviet Union, (Division of Gas in Industrial Purposes and Other Uses)  
12, Bolshaya Konyushennaya Street, Moscow, U.S.S.R., 1979, 227 p. Extent also  
indicated. Original (Russian). Numbered. 5,000 copies printed.  
Lithographed.

Sci. - R. S. Glazunov, Director of Technical Sciences; Zelenin, N. A.; V. I.  
Stepanovskaya; Tikhon, N. A.; Poltina.

Abstract: This collection of articles is intended for specialists engaged in  
developing and operating gas units of industrial enterprises and electric  
power plants.

Contents: The chapter "Gas as a new industrial resource" contains a review of some  
gas uses and methods. Advantages of using natural gas as a source of energy  
are outlined. Different gas burner systems, devices for automatic control  
of the combustion process, structural features of burners operating on natural  
gas, economy of operation and the introduction of new burners. In the  
subsequent chapters and the introduction of new units are described: the basic features  
of gasification and operation of gas units are described. The basic features  
of diagrams of gas-supply systems and equipment. The generalities are  
discussed. One article is followed by references.

CONTENTS:

1. Shchegolev, I. M. Present State and Prospects for Supplying Natural Gas  
to Enterprises and Electric Power Stations With Gas 1
2. The J. Development of Gas Service in the Industry During the  
1955 Period and the Deployment of Natural Gas 2
3. Shchegolev, I. M., A. I. Mal'zeva, A. N. Kuznetsov, and A. T. Reiterberg.  
Methods for Supplying Gas to Industrial Enterprises 8
4. Shchegolev, I. M. Gas Burners for Boilers and Industrial Purposes Which Can  
Be Burned From Gas 21
5. Shchegolev, I. M. Automatic Regulation of the Combustion 69
6. Prokof'yev, I. Yu. Problem of the Economic Profitability of Utilizing Gas  
in Industry 10
7. Shchegolev, I. M. and Yu. A. Kuznetsov. Basic Utilization of Natural Gas 16
8. Shchegolev, I. M. and Yu. A. Kuznetsov. Planning of Future Plants 26
9. Appendix: Library of Congress 26

20-145

(3)

card b/a

DOLOTOV, G.P., inzh.

Safety measures in using natural gas in furnaces and drying apparatus.  
Besop. truda v prom. № no.8:15-18 Ag '59. (MIRA 12:11)

1. Avtomobil'nyy zavod im. Likhacheva.  
(Gas, Natural--Safety measures)

DOLJUV, G.P.

Using natural gas to heat blast air in cupolas. Gaz. prom. 7 no. 5:  
39-41 '62. (MIRA 17:11)

MARIYENBAKH, L.N., doktor tekhn.nauk, prof.; DOLOTOV, G.P., inzh.

Using natural gas for smelting cast iron. Vest.mashinostr. 43  
no.8:65-70 Ag '63. (MIRA 16:9)  
(Cast iron--Metallurgy) (Gas, Natural)

MARIENBAKH, L. M.; DOLOTOV, G. P.

"Die praxis und die perspektiive der anwendung des naturgases bei dem schmelzen  
des gusseisens."

paper submitted for 32nd Intl Cong, Foundry Congress, Warsaw, 13-17 Sep 65.

POLOTOV, G.P., kand. tekhn. nauk; KONDAKOV, Ye.A., inzh.;  
SURINOV, B.P., inzh., retsenzent; FILIMOV, Yu.P.,  
kand. tekhn. nauk, red.

[Design and calculation of industrial furnaces and driers;  
foundry furnaces] Konstruktsiya i raschet zavodskikh pechей  
i ushil' pechi liteinykh tsekhov. Moskva, Mashinostroenie,  
1965. 238 p. (MIRA 18:8)

DARK, S.Ye.; DOLOTOW, G.P.

Smelting cast iron using natural gas. Gas, prov. 10 no. 2:27-31  
165. (MIRA 18:12)

DOLOTOV, I.

Deviation of magnetic compasses. Mor. flot. 24 no.5:  
17-18 My '64. (MIRA 18:12)

1. Starshiy deviator Leningradskoy bazy Okeanrybfloata.

Dolotov, M.

USSR.

UKT-I CUTTER-LOADER FOR THIN SLIGHTLY DIPPING SEAMS. (UGOLNYI KOMBAIN UKT-I Dlya TONIYKH POLOGOPADAYUSHCHIKH PLASTOV). Budryushhev, E. I., Dolotov, M. N. and Savilev, I. P. (Moscow: Ugletekhnidat, 1953, 208pp.; abstr. in Ugol (Coal), Aug. 1954, 48). The design characteristics and operation are described and recommendations are made for organization of work.

KUDRYASHOV, Yevgeniy Ivanovich, DOLOTOV, Mikhail Nikitovich, SAVEL'YEV, Ivan  
Petrovich, SHOKHOVA, A.V., et al.; IONILINA, L.N., tekhn.red.

[UET-2 and UET-2m cutter-loaders] Ugol'nye kombainy UET-2 i UET-2m.  
Moskva, Ugletekhizdat, 1958. 191 p.  
(MIRA 11:8)  
(Coal mining machinery)

DOLOTOV, N. P.

Explosive method of caving in mines of Moshkovogol Combinat. Molotov, N.P.  
and Babokin, I.A., (Ural (Coal). Feb. 1952, 18-21).

The method involves withdrawing some supports and demolishing others.  
It is described in detail. Time and timber are saved in comparison with  
the hand method used previously. (I)

immediate source clipping

DOLOTOV, N.P.; MIKHNOVA, V.N.; SIN'KO, A.I.; BABOKIN, otvetstvennyy red.;  
KOROVENKOVA, Z.A., tekhn. red.

[Handbook for workers and minor grade inspectors in mine transpor-  
tation in the Moscow Basin] Pamiatka dlja rabochikh i mledshego  
nadzora uchastka vnutrisheskogo transporta Podmoskovnogo basseina.  
Moskva, Ugletekhnizdat, 1953. 22 p. (MERA 11#7)  
(Moscow Basin--Mine haulage)

DOLOTOV, Nikolay Pavlovich; RIZNIKOV, G.A., otvetstvennyy redaktor;  
OKHRIMENKO, V.A., redaktor izdatel'stva; NADEINSKAYA, A.A.,  
tekhnicheskiy redaktor

[Manual on longwall work in the Moscow Basin] Pamiatka rabochego  
lavy podmoskovnogo basseina. Moskva, Ugletekhizdat, 1956.  
174 p. (MLRA 10:5)  
(Moscow Basin--Coal mines and mining)

DOLOTOV, N.P.

Industrial training of Moscow Basin miners. Bezop. truda v prom.  
l no.7:6-7 JI '57. (MIRA 10:7)

1. Glavnyy inzhener upravleniya Tul'skogo okruga Gosgortekhnadzora  
SSSR.  
(Moscow Basin--Mining engineering--Safety measures)

"APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000410830003-2

DOLOTOV, N.P., inzh.; ZASLAVSKIY, P.M., inzh.; KLITSUHOB, V.I., inzh.

Observe safety requirements in designing machines and mechanisms.  
Besop. truda v prom. 2 no.7:15-16 J1 '58. (MIRA 11:9)  
(Coal mining machinery)

APPROVED FOR RELEASE: 06/13/2000

CIA-RDP86-00513R000410830003-2"

NECHIFORENKO, M.M., inzh.; DOLOTOV, N.P., inzh.

Safety measures in mines of the Moscow Basin. Bezop.truda. v prom.  
4 no.6:25-27 Je '60.  
(Mira 14:3)

1. Upravleniye Tul'skogo okruga Gosgortekhnadzora RSFSR.

(Moscow Basin—Coal mines and mining—Safety measures)

NECHIPORENKO, M.M.; POLOTOV, N.P., inzh.; SUBBOTIN, A.A., Geroy Sotsialisticheskogo truda; PERMYAKOV, P.N., laureat Leninskoy premii

Effective methods for improving work sanitation in mining. Bezop. truda v prom. 6 no.7:4-6 Jl '62. (MIRA 15:7)

1. Nachal'nik Upravleniya Tul'skogo okruga Gosudarstvennogo komiteta pri Sovete Ministrov RSFSR po nadzoru za bezopasnym vedeniyem rabot v promyshlennosti i gornomu nadzoru (for Nechiporenko). 2. Nachal'nik Tul'skogo kombinata ugol'noy promyshlennosti Podmoskovskogo basseyna Ministerstva ugol'noy promyshlennosti SSSR (for Subbotin). 3. Glavnnyy inzh. Tul'skogo kombinata ugol'noy promyshlennosti Podmoskovskogo basseyna Ministerstva ugol'noy promyshlennosti SSSR (for Permyakov).  
(Tula Province—Coal mines and mining—Safety measures)

SUBBOTIN, A.A., Geroy Sotsialisticheskogo Truda; PERMYAKOV, P.N.,  
laureat Leninskoy premii; NECHIPORENKO, M.M.; DOLOTOV, N.P.

Mechanization and automation in mines of the Priokskiy Economic  
Council. Besop. truda v prom. '7 no.4:2-3 Ap '63.  
(MIRA 16:4)

1. Nachal'nik Tul'skogo kombinata ugol'noy promyshlennosti  
Podmoskovnogo basseyna Ministerstva ugol'noy promyshlennosti  
SSSR (for Subbotin). 2. Glavnnyy inzh. Tul'skogo kombinata  
ugol'noy promyshlennosti Podmoskovnogo basseyna Ministerstva  
ugol'noy promyshlennosti SSSR (for Permyakov). 3. Nachal'nik  
Upravleniya Tul'skogo okruga Gosudarstvennogo komiteta pri  
Sovete Ministrov RSFSR po nadzoru za bezopasnym vedeniyem rabot  
v promyshlennosti i gornomu nadzoru (for Nechiporenko). 4. Glavnnyy inzh.  
Upravleniya Tul'skogo okruga Gosudarstvennogo komiteta  
pri Sovete Ministrov RSFSR po nadzoru za bezopasnym vedeniyem  
rabot v promyshlennosti i gornomu nadzoru (for Dolotov).

(Tula Province--Coal mines and mining)  
. (Automation)

DOLOTOV, N.P.; STARUKHINA, A.D.

Causes of accidents and traumatism at chemical industry enterprises. Bezop. truda v prom. S no.9:22-24 S '64

(MIRA 18:1)

1. Upravleniya Tul'skogo okruga Gosudarstvennogo komiteta pri Srete Ministerov RSFSR po nadzoru za bezopasnym vedeniyem rabot v promyshlennosti i gornomu nadzoru.

NECHIPORENKO, M.M., inzh.; DOLOTOV, N.P., inzh.

Water supply for mine fires in the Moscow Basin. Bezop. truda  
v prom. 8 no.12;16-18 p '64. (MIRA 18:3)

1. Upravleniye Tul'skogo okruga Gosudarstvennogo komiteta pri  
llore Ministerstva RSFSR po nadzoru za bezopasnym vedeniyem  
rabot v promyshlennosti i gornomu nadzoru.

L 24137-66 EWT(m)/EWA(d)/T/EWF(j)/EWP(t) IJP(o) JD/HN/KB/RM  
ACC NR: AP6014665

SOURCE CODE: UR/0314/65/000/008/0040/0041

AUTHOR: Kuzub, V. S. (Candidate of chemical sciences); Dolotova, T. S. (Engineer)

ORG: none

TITLE: Corrosion of metallic materials in the production of diphenylolpropane

SOURCE: Khimicheskoye i neftyanoye mashinostroyeniye, no. 8, 1965, 40

TOPIC TAGS: corrosion, stainless steel, aluminum, titanium, nickel, copper, organic synthetic process, alloy, distillation, corrosion resistant alloy/NIKhMo-20 alloy, 1Kh18N9T stainless steel, Kh18M12M2T stainless steel, Kh21Ni4B stainless steel, Kh23N28M3D3T stainless steel

ABSTRACT: The corrosion resistance of Al (A-DO) (AD-1), Ti (VRL), Ni, Cu alloy NIKhMo-20 and stainless steels 1Kh18N9T, Kh18M12M2T, Kh21Ni4B, Kh23N28M3D3T were studied in the synthesis, distillation and purification processes of diphenylolpropane. The specimens were tested in the following media corresponding to these processes: phenol / BF<sub>3</sub> (2.5%), t = 90-120°C (in the liquid and gaseous phases); diphenylolpropane / CH<sub>3</sub>COOH / impurities, t = 80°C; diphenylolpropane / water / impurities, t = 100°C. For the synthesis and distillation processes the most stable material both in the liquid and gaseous phase is aluminum grade A-DO (grade AD-1 was less stable). Also, the results of tests indicated that an increase in the medium temperature from 90 to 120°C does not affect the corrosion rate of the materials.

Card 1/2

UDC: 620.19:546.621'74:669.15:547.62'213

1 24137-66

ACC NR: AP6014665

studied. Stainless steels and titanium are corrosion-resistant materials in a solution of 30% acetic acid with impurities of phenol, boron fluoride, and resins. In the purification of diphenylolpropane from residues of phenol,  $\text{BF}_3$  and resins containing water vapor, an aqueous solution of phenol and hydrofluoric acid is formed. In this medium copper and stainless steels Kh23N28M3D3T and Kh18Ni2M2T, alloyed with copper and molybdenum and possessing high resistance in hydrofluoric acid, are corrosion-resisting materials. Thus the tests and resulting data permit the recommendation of the metals as structural materials for a specific process in the production of diphenylolpropane. Orig. art. has: 1 table. [JPRS]

SUB CODE: 13, 07 / SUBM DATE: none / ORIG REF: 003

Card 2/2

DOLOTOV, V.A.

Genesis of mountain-steppe fragmentary soils of Kopet Dagh [with summary in English]. Pochvovedenie no. 7:36-41 Jl '58. (MIRA 11:8)

1. Tsentral'nyy Musey pochvovedeniya im. V.V. Dokuchayeva Akademii nauk SSSR, Leningrad.

(Kopet Dagh—Soil formation)

DOLOTOV, V.A.

Agricultural utilization of soils in Vologda Province. Sbor. rab.  
TSentr. nauch. pochv. no.3:192-202 '60. (MIRA 13:9)  
(Vologda Province--Soils)

ZAVALISHIN, A.A.; DOLOTOV, V.A.

In memory of Konstantin Dmitrievich Glinka. Pochvovedenie  
no.9:117-180 S '62. (MIRA 16:1)  
(Glinka, Konstantin Dmitrievich, 1867-1927)

DOLOTOV, V.A.

Fragmental soils as a growth medium for tragacanth-bearing  
milk vetches in Kopet-Dag. Trudy Bot. inst. Ser. 5 no.11:  
212-228 '63. (MIRA 16:10)

DOLOTOV, V.A.

Studying soil formation of arable lands in the east of the Russian Plain. Pochvovedenie no.9: 31-38 Ag [i. e. S] '63. (MIRA 16:10)

1. Tsentral'nyy muzei pochvovedeniya imeni V.V. Dokuchayeva.  
(Kirov Province--Soil formation)  
(Udmurt A. S. S. R.--Soil formation)

DOLOTOV, V.A.

Effect of cultivated plants on soil. Bot. zhur. 48 no.10:1495-1499  
O '63. : (MIRA 17:1)

1. Tsentral'nyy muzey pochvovedeniya imeni V.V.Dokuchayeva, Leningrad.

DOLOTOV, V.A.

Recent data on the liming of acid soils. Priroda 54 no.7:22-23  
Jl '65. (MIRA 18:7)

1. TSentral'nyy muzey pochvovedeniya im. V.V.Dokuchayeva, Leningrad.

DOKHINOV, V.A.

Principles underlying the classification of arable soils.  
Postroyedenie no.315-42 pag 166. (MIL 18:6)

1. Tsentral'nyy musey po pol'zovaniyu imeni V.V.Dokhinyova, Moskva.

DOLCHOV, V.G.; DEMNIKOV, F.Ye.

Adaptability and reliability of an automatic control center.  
Trudy MFI 52:81-84 '63.  
(MIRA 18:3)

L 51857-65 ENT(d)/EWP(v)/T/EWP(x)/EWP(n)/  
ACCESSION NR: AR4041565

EWP(1) Pf-4  
5/02/1/64/001/008/A019/A019  
62.5:658.562

SOURCE: Ref. zh. Avtomat., telemekh. i vycisl. tekhn. Svednyy tom, Abs. 8A133

AUTHOR: Dolotov, V. G.; Temnikov, F. Ye.

TITLE: Adaptability and reliability of a computer

CITED SOURCE: Tr. Mosk. energ. in-ta, vyp. 52, 1963, 95-106

TOPIC TAGS: mass information, information processing, industrial automatic control, supervisory control

TRANSLATION: The results of a development of fundamental units intended for a system of centralized collection and processing mass information are reported. A plant-wide information network encompasses all departmental devices that collect primary information. This network pipes the entire information into a central station where apparatus which determine the arrival and processing of information about the entire process cycle are located. The structure is given. An example of operation is described of a central code-scanning generator (CSG) and a sensor-switching unit. The CSG produces, in the common information network, continuously circulating code scans. The code scans received at the departmental points are used for setting up compensating signals of any physical nature. The central

Card 1/2

L 51857-65

ACCESSION NR: AR4046565

O

sensor-switching unit performs synchronous switching of sensors at all departmental points. The problems of adaptability of the "Tsentrtekhnika-3" centralized information system to the conditions of production are discussed. In such systems, an essential redistribution and degeneration of apparatus, as compared to decentralized and local systems, take place. Only the elements directly associated with the processing, such as sensors and actuators, remain near the production machines; all other apparatus are moved to the center. Reliability of such centers is discussed. Eight illustrations. Bibliography: 1 title.

SUB CODE: DP, II

ENCL: 00

1  
Cord 2/2